



Coli Stop

Indication:

E-Coli, Paratyphoid, Streptococcus, Staphylococcus and bacterial infections in general.

Symptoms:

Loss of appetite, slow crop emptying, loss of weight, vomiting, brown green watery or slimy droppings, sudden death.

Dosage

Prevention: 1 measuring spoon (=5g) for 2 litres drinking water during 1 or 2 days every 2 weeks, preferably starting the day after the race.

As a treatment: 1 measuring spoon (=5g) for 2 litres drinking water during 5-7 successive days.

This cure may be repeated if necessary after a few days.

Coli Stop is a water-soluble powder which can also be given under the feed.

Measuring spoon 1+5 g included.



Advice for use:

- Put the sick animals in quarantine.
- Treat all of your birds, not only the sick ones.
- Mix the needed dose in an amount of drinking water or feed the animals consume within one day.
- Be sure no other water is available during the treatment
- After medication give your animals MULTIVITAMIN (the day after last day of treatment).
- Give a maintenance dose of MULTIVITAMIN every week.
- Disinfect lofts/houses.
- Do not give the day of shipping.
- In case of doubt or persistent problems, consult a vet. surgeon.

NOTES: READ INSTRUCTIONS BEFORE USE.

- Store dry, out of reach of direct sunlight, between 0-30 °C.
- Shelf life after first opening; One (1) year.
- Batchnr. and expiry date: see package.
- Avoid contact of the medicine with your hands.
- Keep out of reach of children.
- Interaction(s): not known. Contra indication(s) overdosage.
- For animal treatment only. Not for animals intended to produce food for human consumption.
- Dispose of empty packaging and remaining product in the household refuse.
- This veterinary medicine is marketed in accordance with the small animal exemption scheme.

Contains:

Streptomycine Sulfate 13,9%, Neomycine Sulfate 4%, Colistine Sulfate 2%. Inactive ingredients q.s. to 100 g.
