



# General Care

---

## Indication:

Canker (trichomoniasis, hexamitosis), paratyphoid, E-coli and bacterial infections in general.

## Symptoms:

Loss of weight and general condition, the droppings are slimy and green, legs and wings joints are swollen, eggs are infertile, they sometimes twist their neck on the side, Young bird sickness, sudden death.

## Dosage

Prevention: 1 measuring spoon (=5g) for 2 litres drinking water during 1 or 2 days every 2 weeks, preferably starting the day after the race.

As a treatment: 1 measuring spoon (=5g) for 2 litres drinking water during 5-7 successive days.

This cure may be repeated if necessary after a few days. Stir well before use!

General Care is a water-soluble powder which can also be given under the feed.

Measuring spoon 1+5 g included.



## Advice for use:

- Put the sick animals in quarantine.
- Treat all of your birds, not only the sick ones.
- Mix the needed dose in an amount of drinking water or feed the animals consume within one day.
- Be sure no other water is available during the treatment.
- Do not provide any calcium, grit, salt or dairy products during treatment.
- After medication give your animals MULTIVITAMIN (the day after last day of treatment).
- Give a maintenance dose of MULTIVITAMIN every week.
- Disinfect lofts/houses.
- Do not give the day of shipping.
- In case of doubt or persistent problems, consult a vet. surgeon.

## NOTES: READ INSTRUCTIONS BEFORE USE.

- Store dry, out of reach of direct sunlight, between 0-30 °C.
- Shelf life after first opening; One (1) year.
- Batchnr. and expiry date: see package.
- Avoid contact of the medicine with your hands.
- Keep out of reach of children.
- Interaction(s): not known. Contra indication(s) administration with Bromhexine and overdosage.
- For animal treatment only. Not for animals intended to produce food for human consumption.
- Dispose of empty packaging and remaining product in the household refuse.
- This veterinary medicine is marketed in accordance with the small animal exemption scheme.

## Contains:

Furaltadone HCl 12%, Metronidazole 20%, Chlortetracycline HCl 10%, Inactive ingredients q.s. to 100 g.

---