



Intestinal

Indication:

Adeno-Coli Syndrome, Coccidiosis, Canker and bacterial infections in general.

This mix contains a combination of active components that are effective against the Adeno/E-coli syndrome and Coccidiosis. The Adeno/E-coli syndrome is a problem, that is found in young pigeon populations and can be very harmful. Early treatment is important.

Symptoms:

Loss of weight and general condition, vomiting, the droppings are slimy and green, death.

Dosage

Prevention: 1 measuring spoon (=5g) for 2 litres drinking water during 1 or 2 days every 2 weeks, preferably starting the day after the race.

As a treatment: 1 measuring spoon (=5g) for 2 litres drinking water during 5-7 successive days. Stir well before use!

This cure may be repeated if necessary after a few days.

Intestinal is a water-soluble powder which can also be given under the feed.

Measuring spoon 1+5 g included.



Advice for use:

- Put the sick animals in quarantine.
- Treat all of your birds, not only the sick ones.
- Mix the needed dose in an amount of drinking water or feed the animals consume within one day.
- Be sure no other water is available during the treatment.
- After medication give your animals MULTIVITAMIN (the day after last day of treatment).
- Give a maintenance dose of MULTIVITAMIN every week.
- Disinfect lofts/houses.
- Do not give the day of shipping.
- In case of doubt or persistent problems, consult a vet. surgeon.

NOTES: READ INSTRUCTIONS BEFORE USE.

- Store dry, out of reach of direct sunlight, between 0-30 °C
- Shelf life after first opening; One (1) year.
- Batchnr. and expiry date: see package.
- Avoid contact of the medicine with your hands.
- Keep out of reach of children.
- Interaction(s): not known. Contra indication(s) overdose
- For animal treatment only. Not for animals intended to produce food for human consumption
- Dispose of empty packaging and remaining product in the household refuse
- This veterinary medicine is marketed in accordance with the small animal exemption scheme.

Contains:

Trimethoprim 2,5%, Sulfadiazine Sodium 12,5% , Furaltadone HCl 5%, Ronidazole 5% , Inactive ingredients q.s. to 100 g.