

Salmo-Stop ^{Extra Strong}

Salmonella (paratyphoid), E-coli and bacterial infections

Symptoms

Diarrhea (green diarrhea, in some vomiting birds), slow curvature of the neck, infectious arthritis (limp or "fallen" wing), subcutaneous abscesses (usually under the eyes, in the form of small, yellow and hard purulent sacs), infertility, sudden death.



100 g POWDER

Dosage

Prevention: 1 measuring spoon (=5g) for 2 litres drinking water during 1 or 2 days every 2 weeks, preferably starting the day after the race.

As a treatment: 1 measuring spoon (=5g) for 2 litres drinking water during 5-7 successive days. This cure may be repeated if necessary after a few days.

Contains: Furaltadone HCl 12,5%, Chlortetracycline HCl 12,5%, Chloramphenicol 8%. Inactive ingredients q.s. to 100 g

50 pcs TABS

Dosage

Preventive: 1 tablet per pigeon (400-500 g bodyweight) for 1-2 consecutive days.

Curative: 1 tablet per pigeon (400-500 g bodyweight) for 3-5 consecutive days, or as Vet. prescribes.

Contains: Chlortetracycline HCl 78 mg/g ; Furaltadone HCl 80 mg/g ; Chloramphenicol 50 mg/g