



Moulting season

"A good moult is the basis for optimal results next year"



In case of any suspicion of Paratyphoid (Salmonellosis) also called "leg" or "wing" illness, one is recommended to cure the pigeons for 10 to 14 consecutive days with **General Care 4 in 1** (C4B013), (5 g per 2 litres of water), followed by 2 days **Caremin** (C4B049) / 10 ml per 1 litre of drinking water/ or **Multivitamin**(C4B032) /5 g per 2 litres of water or 5 g per 1 kilogram of grains/*

* Grain can be moistened with **Carechol**(C4B048)

In case of this illness in the surrounding area of your loft we advise you to treat for 7 consecutive days with **Salmo Stop** (C4B019) (5 g per 2 litres of water) preventive, followed by 2 days **Multivitamin**(C4B032) (5 g per 1 litre of water or 5 g per 0,5 kg kilogram of grains (*))

* Grain can be moistened with **Carechol**(C4B048)

At the beginning of the moulting season 1 consecutive week **Caremin** (C4B049) / 10 ml per 1 litre of drinking water/ or **Multivitamin**(C4B032) /5 g per 2 litres of water or 5 g per 1 kilogram of grains/

* Grain can be moistened with **Carechol**(C4B048)

Followed by once or twice per week **Carechol**(C4B048) (10 ml per 1 l. water).
Caremin (C4B049) .

Cure the pigeons 5 consecutive days with **Combi Mix 4 in 1**(C4B011) (5 g per 2 litres of water).

Once a week **Caremin** (C4B049) / 10 ml per 1 litre of drinking water/ or **Multivitamin**(C4B032) /5 g per 2 litres of water or 5 g per 1 kilogram of grains/*

* Grain can be moistened with **Carechol**(C4B048).

Tip! We advise you to check the droppings of your pigeons after the racing season for worms, coccidiosis and paratyphoid. This check has to be done before the start of the moulting period.